

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 April 2001 (19.04.2001)

PCT

(10) International Publication Number
WO 01/26649 A1

- (51) International Patent Classification: **A61K 31/205, 31/22, A61P 13/12**
- (21) International Application Number: **PCT/IT99/00317**
- (22) International Filing Date: **11 October 1999 (11.10.1999)**
- (25) Filing Language: **Italian**
- (26) Publication Language: **English**
- (71) Applicant (for all designated States except US): **SIGMA-TAU INDUSTRIE FARMACEUTICHE RIUNITE S.p.A. [IT/IT]; Viale Shakespeare, 47, I-00144 Rome (IT).**
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): **ARDUINI, Arduino [IT/IT]; Via della Camilluccia, 701, I-00135 Rome (IT).**
- (81) Designated States (national): **AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**
- (84) Designated States (regional): **ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).**
- Published:
— With international search report.
- (74) Agent: **SPADARO, Marco; Sigma-Tau Industrie Farmaceutiche Riunite S.p.A., Viale Shakespeare, 47, I-00144 Rome (IT).**
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/26649 A1

(54) Title: **USE OF L-CARNITINE AND ITS ALKANOYL DERIVATIVES AS OSMOTIC AGENTS IN SOLUTIONS FOR MEDICAL USE**

(57) Abstract: **The use of L-carnitine and its alkanoyl derivatives, optionally in the form of a pharmaceutically acceptable salt, as osmotic agents in the preparation of solutions for medical use, particularly for peritoneal dialysis, is described.**

"Use of L-carnitine and its alkanoyl derivatives as osmotic agents in solutions for medical use".

The invention described herein relates to the use of L-carnitine and its alkanoyl derivatives, optionally in the form of a pharmaceutically acceptable salt as osmotic agents in solutions for medical use, particularly in peritoneal dialysis.

Background of the invention

Patients suffering from end-stage renal disease (or ESRD) must either undergo dialysis therapy or be submitted to a kidney transplant. Both therapeutic interventions are extremely demanding, both from the point of view of the quality of life of the patient and in terms of social costs. For a review of dialytic therapy see, for example, Pastan S. and Bailey J. in the New England Journal of Medicine, 14 May 1998, pp. 1428-1436, incorporated herein for reference in its entirety.

Dialytic therapy comprises two types of treatment, namely peritoneal dialysis and haemodialysis. There are major differences between the two types of dialysis, such as, in the case of haemodialysis, the cost of the therapy, the need for dedicated departments with expensive equipment and qualified staff, and the quality of life of the patient. Peritoneal dialysis, on the other hand, enjoys greater favour on account of its simplicity of execution, which can be handled by the patient himself in the form of self-medication. In Italy, for example, 15% of dialysis patients use peritoneal dialysis, which is practically the same as in the USA (16%), while the percentages of patients on peritoneal dialysis are higher in Canada (38%) and the United Kingdom (52%), and get up to as much as 90% in Mexico. The reason for these different rates is also to be

./.

attributed to the lower cost of peritoneal dialysis as compared to the cost of haemodialysis which not all national health systems are prepared to bear. We should, however, not overlook the fact that peritoneal dialysis allows the patient to maintain a less
5 constrained life-style, since the dialysis session can be planned with a certain measure of autonomy in the course of the person's normal activities. In addition, automatic devices also allow dialysis during the hours of the night.

Despite this, the choice between the two types of dialysis
10 is not a free one; for instance, peritoneal dialysis is indicated for patients with cardiac insufficiency or unstable angina who cannot support the alterations of blood flow and/or arterial blood pressure that accompany the haemodialysis session (see reference cited above).

15 One can postulate a therapeutic progression for the ESRD patient which starts with peritoneal dialysis, proceeds via haemodialysis and finally reaches a stage where a kidney transplant is needed.

Peritoneal dialysis is not without disadvantages and
20 unwanted adverse effects. These drawbacks can be placed in two distinct, even if related, categories, namely adverse clinical effects and technological problems. The purpose of the invention described herein is to remedy these disadvantages and adverse effects.

25 In the typical execution of a peritoneal dialysis session, a plastic catheter is implanted in the peritoneal cavity and anchored to the subcutaneous tissues. A dialysis solution contains physiological amounts of sodium, calcium, magnesium, compatible physiological buffer and a non-toxic
30 osmotic agent, of such a nature as to make the solution

./.

hyperosmolar as compared to the plasma. The solution is infused via the catheter into the peritoneal cavity where it then remains for several hours. During this time, the peritoneal membrane exchanges solutes by diffusion in such a way as to obtain replacement with fresh fluid. Given that renal function decreases in the first few years of dialysis, the dose of dialysis fluid to be exchanged increases in the course of time.

Peritonitis is the serious complication that occurs most frequently. Other types of complications are losses of amino acids and albumin, incompatibility of the dialysis solution, volume effects in the peritoneal cavity, metabolic consequences, symptoms affecting the digestive tract, reduced appetite and others (for a review see C.M. Mion, R. and Gokal and N.P. Mallick, Lancet, 1999; 353; 823-28).

One of the most pressing problems in the peritoneal dialysis sector is the choice of a suitable osmotic agent.

The requisites of an ideal solution for peritoneal dialysis include:

- supplying the nutritional requirement and avoiding adverse metabolic effects;
- ensuring minimal absorption of the osmotic agent, which, in any case, must be non-toxic;
- being capable of correcting acidosis and having a physiological pH;
- in addition to considerations with technological implications, such as apyrogenicity, absence of metals and residues of synthetic materials, the solution must also inhibit bacterial and fungal growth, must not damage the immune defences and must be inert in relation to the peritoneal membrane.

A typical solution for peritoneal dialysis contains glucose in various concentrations as an osmotic agent, and various amounts of lactate (which has replaced acetate owing to problems of intolerance on the part of the patient), sodium, potassium and calcium. Buffer systems have also been studied in an attempt to solve the problem of sterilization and stabilization of the solution.

As regards the sterilization aspect, this is a critical technological problem; in fact, heat sterilization, commonly used in the sector of solutions for medical use, causes degradation of glucose, with consequent production of toxic secondary derivatives, such as aldehydes and 5-hydroxy-methylfurfural. Traditionally heat sterilization of the solution containing glucose (also indicated as dextrose) is done at a pH between 5.0 and 5.5, precisely in order to avoid caramelization of the glucose. The acid pH leads to further problems for the patient using the solution, such as, for example, abdominal pain and sclerosis of the peritoneal membrane, which entails a reduction of the elimination of solutes (Schmidt et al., Arch. Int. Med., 141; 1265-1266, 1981).

The purpose of the invention described herein is also to provide a solution to the complex problems related to the use of glucose as an osmotic agent in solutions for peritoneal dialysis.

Glucose is extensively used owing to its great availability on the market and its low cost. It is a relatively safe substance, but its use at high concentrations and its prompt absorption lead to short ultrafiltration times, and metabolic complications, such as hyperinsulinaemia, hyperlipidaemia, and weight gain. In addition, hyperosmolarity and low pH can damage the mesothelium and macrophages. Moreover, the potential

./.

glycosylation of stromal proteins leads to further damage to the peritoneum. Also reported is the inhibition of phagocytosis, bactericidal activity and the synthesis of LTB₄ in peripheral blood neutrophils. In continuous ambulatorial peritoneal dialysis (CAPD), where the application time can be as much as 6 hours or more, the glucose concentrations are very high to be able to maintain the ultrafiltration capacity. For a review of the biocompatibility of solutions for peritoneal dialysis see C.J. Holmes in Peritoneal Dialysis International, Vol. 13, pp. 88-94, 1993.

To overcome the problems created by the use of glucose as an osmotic agent in peritoneal dialysis, the state of the art directs experts in the field towards two distinct types of solution:

- 1) the use of low-molecular-weight osmotic agents, capable of sustaining ultrafiltration with minimal metabolic effects, without, however, altering the ultrafiltration profile;
- 2) the use of high-molecular-weight osmotic agents in an attempt to act on both factors.

Of the various low-molecular-weight agents proposed, to date only glycerol and mixtures of amino acids would appear to be of a certain clinical interest. In Italy, for example, a 1.1% multi-amino-acid solution is being marketed by Baxter under the trade mark Nutrineal® PD2 and PD4.

These proposed alternatives to glucose are not problem-free; other saccharides have metabolic effects: for example, fructose gives rise to hypertriglyceridaemia and hyperosmolarity, sorbitol hyperosmolarity and accumulation, xylitol lactic acidosis and hyperosmolarity; glycerol is well tolerated, but its ultrafiltration capacity is short-lasting and it also causes hyperosmolarity,

./.

while an adverse effect on phagocytes has also been reported (de Fijter CWH et al., *Advances in Continuous Ambulatory Peritoneal Dialysis*, Toronto, Peritoneal Dialysis Publication, 1991, 154-7). Amino acids, though useful in undernourished patients, give rise to acidosis and to an increase in the nitrogen load, which is contraindicated in an uraemic patient. On the other hand, high-molecular-weight osmotic agents present a whole series of disadvantages of their own, such as possible immunogenicity in the case of peptides, absorption, intraperitoneal bleeding (demonstrated in rats) and ultrafiltration in the case of dextrans (MW 60-200 kDa), cardiovascular instability, peritoneal damage and haemorrhage in the case of polyanions and cations (MW 40-90 kDa), prolonged half-life, immunogenicity, allergenicity and high viscosity of the solution in the case of gelatines (MW 20-390 kDa) and maltose retention in the case of glucose polymers.

Unfortunately, the adverse effects of solutions for peritoneal dialysis do not stem only from the osmotic agent chosen, but also from other components of the solutions. Lactate, for example, when combined with the low pH of the solution, which is necessary in order to be able to perform sterilization, suppresses various functional activities of the peripheral and peritoneal leukocytes and inhibits the production of IL-6 and TNF α by mononuclear cells.

In his review of osmotic agents, Gokal concluded in 1990 that at that time there was no osmotic agent capable of replacing glucose (Coles GA, Davies M, Williams JD (eds): *CAPD: Host Defence, Nutrition and Ultrafiltration*. Contrib. Nephrol., Basel, Karger, 1990, vol. 85, pp. 126-133).

An enormous effort is being made to find an alternative osmotic agent to glucose, meeting or at least coming close to meeting the requisites of the "ideal" solution. Among the numerous patent references, we should cite patent JP 11071286, filed in the name of the Terumo Corp., which describes a solution where the osmotic agent consists in a mixture of glucose and maltose in molar ratios of 1:0.05-5 and with an osmotic pressure of 280-600 mOsm/kg at pH 6.0-7.5, with enhanced water removal characteristics and reduced glucose absorption. For obese, diabetic patients, the same company supplies a complex osmotic agent, made up of N-acetylamine acid (L-amino acid), N-acetyl-D-glucosamine, glucuronic and/or ascorbic acid (patent JP 11071273). Mixtures of saccharides with hexavalent alicyclic alcohols, hexonic acid and sacchric acid are described in patent JP 11049671, filed in the name of Baxter Int. Inc. Patent application WO 9901144, filed by Allied Therapeutics Ltd., describes synthetic hydrogenated di- and trisaccharides. Patent MX 9601855, filed by Trevino, uses dextran 60. Baxter again, in patent JP 10094598, proposed non-reducing oligosaccharides or polysaccharides containing from 3 to 12 residues. In patent application WO 9801141, filed by Bieffe Medital SpA, the use of glycosaminoglycans, devoid of anticoagulant or haemorrhagic activity, is described. US patents US 5629025, US 5589197 and US 5631025, filed in the name of Baxter International Inc., describe solutions for peritoneal dialysis with a low sodium content, for which substances containing at least one amino acid or polypeptide, or a polyglucose, are envisaged as osmotic agents. The University of Missouri supplies chemically crosslinked gelatine as an osmotic agent to partially or wholly

./.

replace glucose (US 4604379). Starch hydrolysates are described in US 5837060, filed in the name of Roquette Freres. Patent JP 7323084, filed by Morishita Roussel KK and Ajinomoto Co. Inc., describes the use of trehalose to prepare
5 neutral solutions to replace glucose. See also US patent US 4761237.

Examining in greater detail the low-molecular-weight osmotic agents to which the present invention refers, the state of the art provides teachings directed to the use of amino acids
10 or short peptides, which are advantageous from the point of view of the nutritional support of undernourished patients. Baxter International Inc., under its US patent US 5776503, supplies a mixture of amino acids, which is highly complex but, despite its very high cost, has yet to be surpassed by the many
15 alternatives proposed. US patent US 5780438, filed in the name of Giltech Limited, describes a stable solution, where the osmotic agent consists of a mixture of peptides obtained from the proteolysis of casein or whey proteins. US patent US 5869444, filed in the name of Research Corporation
20 Technologies, extensively discusses the alternatives to glucose and directs the experts in the field towards low-molecular-weight osmotic agents of an amino-acid nature. However, though admitting the nutritional benefit, mentioned earlier, the inventors stress the disadvantage of the high cost, and the
25 increased nitrogen load in the blood. Thus, in the cited patent, they propose the use of oligopeptides (300-2000 Da) derived from the enzymatic hydrolysis of high-quality proteins, such as whey, which are advantageous both from the functional point of view of the dialysis and from the nutritional point of view. In
30 this patent, however, the need is perceived for a very thorough

./.

and carefully controlled hydrolysis and separation process in order to avoid the risk of high-molecular-weight components, potential antigens or allergens. Among the protein sources, those mentioned include collagen, the use of which today is, however, questionable owing to problems of prion contamination (BSE, scrapie), milk proteins, but not casein, and others. In the course of the description, the inventors admit a series of difficulties in assuring the quality of the hydrolysis process.

In relation to other aspects of peritoneal dialysis, DE 19748290, WO 991762, JP 10330270, WO 9852599, WO 9850060, CA 2219822, WO 9917762 and US 5827820 are cited.

All documents cited are incorporated herein for reference in their entirety.

Abstract of the invention

It has surprisingly been found that L-carnitine, or, alternatively, one of its lower alkanoyl derivatives, or L-carnitine in combination with its lower alkanoyl derivatives, is useful as an osmotic agent in the preparation of solutions for peritoneal dialysis, and in general as an osmotic agent for solutions for medical use.

One subject of the invention described herein is the use of L-carnitine and its lower alkanoyl derivatives, where what is meant by lower alkanoyl derivative is a straight or branched aliphatic acyclic residue with from 2 to 8 carbon atoms, optionally in the form of a pharmaceutically acceptable salt, as an osmotic agent in solutions for medical use, particularly for the preparation of solutions for peritoneal dialysis. Another subject of the present invention consists in solutions for medical use characterised in that the osmotic agent is L-carnitine or one of its alkanoyl derivatives, as defined above, optionally in

./.

combination with one another or with one or more known osmotic agents.

L-carnitine and its lower alkanoyl derivatives are known to have various therapeutic uses. In particular, US patent US 4327167, filed in the name of the applicant, describes the use of alkanoyl carnitines, as defined above, in a therapeutic method for the treatment of chronic uraemic patients undergoing regular dialysis. Also described are polysaline solutions for haemodialysis containing an alkanoyl carnitine. Patent EP 0793962, filed in the name of the applicant, describes the use of propionyl L-carnitine for the preparation of a medicine useful for the selective treatment of chronic obliterating atherosclerosis (claudicatio intermittens). Patent IT 1155772, filed in the name of the applicant, describes the use of alkanoyl L-carnitine in the therapy of myocardial anoxia, ischaemia, arrhythmia syndromes and heart failure. Patent US 4255449, filed in the name of the applicant, describes the use of L-carnitine in the treatment of dyslipidaemias. Patent application WO 9906039, filed in the name of the applicant, describes the use L-carnitine and its alkanoyl derivatives in combination with polycosanols for the treatment of dyslipidaemias. There are numerous descriptions of combinations of L-carnitine and alkanoyl derivatives with other active ingredients, e.g. gamma-linoleic acid (see WO 9841113) for the treatment and prevention of the side effects of diabetes mellitus, particularly peripheral neuropathy.

US patent US 4272549 teaches the use of particular administration regimens of L-carnitine combined orally and intravenously to combat post-dialysis syndrome in uraemic patients undergoing regular haemodialysis treatment.

US patent US 4237167 teaches the use of particular administration regimens of acyl L-carnitine combined orally and intravenously to combat post-dialysis syndrome in uraemic patients undergoing regular haemodialysis treatment.

5 Patent application WO 9907419, filed in the name of Gupta, describes compositions for dialysis containing an effective amount of at least one vitamin selected from the group consisting of folic acid, vitamin B₆, thiamine, vitamin B₁₂, and optionally vitamin C and/or carnitine. The purpose of these
10 preparations is to compensate for the loss of vitamins to which patients on either haemodialysis or peritoneal dialysis are subject. The effective amounts are indicated in the description. In the case of free L-carnitine, an amount less than 50 $\mu\text{mol/l}$ is specified for the dialysed patient during each dialysis session for
15 the prevention of vitamin and carnitine deficiencies. The preferred concentrations range from 50 to 300 $\mu\text{mol/l}$. Thus, the amount of L-carnitine present in the solutions described by Gupta is less than the amount necessary for L-carnitine to act as an osmotic agent.

20 The advantages of using L-carnitine or one of its alkanoyl derivatives, as defined above, are multiple. The replacement of glucose by L-carnitine or one of its alkanoyl derivatives eliminates the adverse effects described above. Furthermore, the carnitines (meaning L-carnitine or its alkanoyl derivatives as
25 defined in the invention described herein) are compatible with the bicarbonate buffer, and therefore obviate the disadvantage of having to use solutions at below physiological pH, such as the pH 5.0 or 5.5 typical of glucose solutions.

The use of carnitines, particularly L-carnitine, acetyl L-carnitine and propionyl L-carnitine, is additionally advantage-
30

./.

ous compared to other known osmotic agents because the carnitines are non-toxic, well tolerated substances with no adverse effects at the doses described below. Unlike the amino acids, the carnitines play no part in protein metabolism and thus do not aggravate the nitrogen load of the uraemic patient. As regards the difference from high-molecular-weight osmotic agents, the advantage is immediate: the carnitines are natural substances present in living organisms, particularly mammals, including man. For this reason, the risk of introducing xenobiotic substances into the body is eliminated.

Moreover, the use of carnitines as osmotic agents provides the dialysed patient with the amount of L-carnitine necessary to compensate for the carnitine losses which occur during the dialysis session. For data on carnitine levels in patients undergoing CAPD, see Kidney Int. 1996 Jan; 49(1): 158-62 and Perit. Dial. Int. 1993; 13 Suppl 2.

A further advantage of the application of the invention described herein is that carnitine, when used as an osmotic agent, not only compensates for the carnitine losses, but is also capable of exerting its own therapeutic effects in a series of diseases related to renal insufficiency, such as, for example, the diseases described in the above-mentioned patents.

The invention will now be described in detail also with the aid of examples. Further subjects of the invention described herein, with their respective advantages, will be apparent to experts in the field to which the present invention pertains.

Detailed description of the invention

What is meant by lower alkanoyl is an acyl group with from 2 to 8 carbon atoms, preferably from 2 to 6, such as acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, 2-methyl-

./.

butyryl, 2,2-dimethylpropionyl, hexanoyl, heptanoyl, octanoyl and all their possible isomers.

The invention described herein envisages the use of carnitines as inner salt. If deemed suitable, one of their
5 pharmaceutically acceptable salts can be used. What is meant by pharmaceutically acceptable salt of L-carnitine or of an alkanoyl L-carnitine derivative is any salt of the latter with an acid which does not give rise to unwanted toxic or side effects. These acids are well known to pharmacologists and to experts in
10 pharmaceutical technology.

Examples of pharmaceutically acceptable salts of L-carnitine or alkanoyl L-carnitines, though not exclusively these, are chloride, bromide, orotate, acid aspartate, acid citrate, acid phosphate, fumarate and acid fumarate, maleate and acid
15 maleate, acid oxalate, acid sulphate, glucose phosphate, tartrate and acid tartrate. The preferred salts are those with fumarate, aspartate, citrate and maleate.

Also the subject of the invention described herein are solutions for peritoneal dialysis, both in the form of ready-to-use
20 solutions and in the form of concentrates to be diluted at the time of use, containing an osmotic agent according to the present invention.

The dosages, posology and treatment regimen in general will be determined by the primary care physician according to
25 his knowledge of the case, the patient's condition and the extent of the disease to be treated.

In a first preferred realization, in the invention described herein L-carnitine, inner salt, is used.

In a second preferred realization, carnitine is present in
30 the form of a salt with fumaric acid. Though not wishing to be

./.

bound by any theory, the applicant believes that the fumarate salt may be particularly advantageous for supplying the energy requirement of the dialyzed patient. Fumarate, in fact, is an energy substrate which is useful in the treatment of organ ischaemia. The applicant has demonstrated the efficacy of L-carnitine fumarate in the treatment of organ ischaemia, particularly ischaemic heart disease, as described in patent application 99RM0003328, incorporated herein for reference in its entirety.

10 In a third realization of the invention described herein, a combination of L-carnitine and acetyl L-carnitine is used. This combination is additionally advantageous to provide the patient with an acetyl L-carnitine supplement.

In the description of the possible realizations of the invention, what is meant by carnitine is L-carnitine, as inner salt, or a salt with a pharmaceutically acceptable acid, as described above, alone or in combination with one of its alkanoyl derivatives as inner salt, or as a salt with a pharmaceutically acceptable acid, or one of its alkanoyl derivatives, as inner salt or a salt with a pharmaceutically acceptable acid.

20 In a first aspect of the present invention, carnitine is used as an osmotic agent as a total substitute for glucose.

The carnitine concentrations are those sufficient for it to act as an osmotic agent and concentrations up to a physiologically tolerable maximum are envisaged. It is understood that the carnitine concentration will be such as to ensure a satisfactory effect for the uses envisaged for the present invention. In particular, it is considered satisfactory the

obtained which can be considered a therapeutic effect in the context of peritoneal dialysis.

Where not otherwise specified, the concentrations are understood to be weight/volume (w/v).

5 Examples of concentrations are approximately 0.5 to approximately 10%, preferably approximately 0.7 to approximately 7%, and more preferably from approximately 1 to 5%. In a typical realization of the present invention, the carnitine concentrations are those normally used for glucose in
10 commercial preparations, namely from 1.5 to 4.25%.

It is understood that experts in the field will be capable of determining effective concentrations according to the type of solution used. Examples are concentrations starting from approximately 0.5%.

15 If so required, alternatively, carnitine can be used as a partial substitute for glucose. The respective carnitine and glucose concentrations can be freely varied, provided that a satisfactory effect is obtained in terms of the uses envisaged for the present invention. Examples of combinations with glucose
20 are 4.0% glucose - 0.25% carnitine; 1.0% glucose - 0.5% carnitine; 0.5% glucose - 1.0% carnitine; 0.25% glucose - 4.0% carnitine. The 0.5% glucose - 1.0% carnitine combination is preferred.

Other possible realizations of the invention consist in a
25 combination of carnitine as an osmotic agent with other known osmotic agents; for example, preferred combinations are those with amino acids, such as the formulations already present on the market, or with the dipeptides and/or polypeptides of the above-mentioned patents. One particularly advantageous
30 realization is the use of carnitine in the twin bags described in

./.

patent DE 19748290, which uses bicarbonate buffer. It is also useful to increase the dose of carnitine, particularly L-carnitine, in the solutions described in WO 9907419 up to a concentration which is effective as an osmotic agent.

5 In another possible realization, the osmotic agent according to the present invention is used in combination with the osmotic agents described in US patent US 5827820, filed in the name of Baxter International Inc.

The use of the osmotic agent according to the present
10 invention is also envisaged in combination with high-molecular-weight osmotic agents, such as, for instance, those described in the above-cited references, and particularly with icodextrin.

In a particular realization of the present invention, a surfactant normally used in this field is added to the peritoneal
15 dialysis solution. Specific mention is made of palmitoyl L-carnitine.

One specific subject of the present invention consists in solutions for medical use characterized in that the osmotic agent is L-carnitine and/or its alkanoyl derivatives, in which the
20 alkanoyl is a straight or branched aliphatic residue with from 2 to 8 carbon atoms, optionally in the form of a pharmaceutically acceptable salt. A particular subject of the present invention consists in solutions for peritoneal dialysis.

As regards those aspects pertaining to industrial
25 applicability, the solutions which are the subjects of the present invention will be contained in suitable containers for peritoneal dialysis, generally bags made of suitable material compatible with medical use. Containers for peritoneal dialysis are known to experts in the field and do not require any particular
30 description, the reader being referred to the specific literature

./.

and to the general knowledge of the technical field to which the invention pertains. Examples are bags with a single chamber or multiple chambers, e.g. a double chamber, or separate bags containing different solutions to be mixed at the time of use by means of automatic equipment. Containers for peritoneal dialysis containing a solution according to the present invention are covered by the protection afforded by the present patent application.

The invention described herein is now described with the aid of experimental tests permitting the implementation of the the realization preferred. It is fully understood that equivalent realizations coming within the framework of the present invention can be implemented by the person having ordinary skill in the field, availing himself only of his own general knowledge, even by the trial and error method, without any need for further description on the part of the present invention.

***In-vitro* transport studies**

Fluid transport *in vitro* was performed using tubes consisting of semipermeable cellulose membranes containing the various different dialysis solutions.

Buffer solutions were added with scalar concentrations of carnitine (0.5, 1.0 and 1.5%) in bicarbonate buffer (30 mM) and NaCl (100 mM) at pH 7.2. A 1.5% glucose solution was used as a reference.

The composition of the buffer solution is as follows: sodium 134 mmol/l, calcium 1.75 mmol/l, magnesium 0.5 mmol/l. The solutions containing glucose were buffered at pH 5.5 with 35 mmol/l of L-lactate. The solutions containing carnitine were buffered as for glucose at pH 7.0-7.6 with 34 mmol/l of bicarbonate. Ten ml of the various dialysis solutions

./.

were placed in the tubes and the tubes were suspended in a 1-litre graduated cylinder filled with a 0.9% NaCl solution. The saline bath was recycled at a rate of 500 ml/min with direct flow along the main axis of the dialysis tube using an infusion pump.

- 5 The amount of fluid recovered inside the tube was determined gravimetrically after removing the fluid adhering to the walls of the membrane with a sheet of absorbent paper. The tube was then put back inside the cylinder and submitted to successive weight measurements at 15, 30, 45, 60, 90, 120, 180, 240, 300
10 and 360 minutes.

Table 1 gives the increase in weight of the fluid recovered from the dialysis solution in the course of time.

TABLE 1.

- 15 Fluid transport induced by 10 ml of fluid containing different concentrations of carnitine or glucose.

Times (min)	Carnitine 0.5% (g fluid)	Carnitine 1.0% (g fluid)	Carnitine 1.5% (g fluid)	Glucose 1.5% (g fluid)
15	0.02±0.01	0.07±0.02	0.10±0.01	0.09±0.02
30	0.06±0.01	0.13±0.02	0.26±0.03	0.19±0.03
45	0.10±0.03	0.20±0.03	0.30±0.02	0.27±0.02
60	0.12±0.02	0.22±0.03	0.35±0.04	0.31±0.04
90	0.17±0.03	0.33±0.03	0.48±0.04	0.41±0.04
120	0.19±0.02	0.36±0.03	0.54±0.04	0.47±0.04
180	0.21±0.03	0.42±0.04	0.66±0.04	0.54±0.04
240	0.27±0.02	0.49±0.03	0.77±0.05	0.59±0.04
300	0.29±0.02	0.51±0.03	0.77±0.04	0.63±0.05
360	0.27±0.03	0.48±0.04	0.74±0.05	0.62±0.04

./.

The values are means ($n=3$) \pm S.D. of 3 different experiments.

The weight of the dialysis tube increases progressively over time as a function of the different carnitine concentrations. The plateau is reached for all concentrations assayed at 240 minutes. The trend of the samples containing 1.5% glucose is comparable to that of the samples containing 1.5% carnitine.

***In-vivo* experiments**

The peritoneal dialysis experiment was conducted in male Sprague-Dawley rats weighing 500-600 g (Charles River) maintained on a standard diet with water *ad libitum*. The animals were anaesthetized with an intraperitoneal injection of inactin (100 mg/kg) and were placed on an operating table at controlled temperature. The animals were submitted to a tracheostomy to cannulate the left jugular vein with a PE50 medical silicone tube. Thirty minutes after administration of the anaesthetic the animals were infused with a saline solution at a rate of 2.3 ml/h throughout the period of the experiment. The dialysis solution (15 ml), after preheating to 37°C, was inoculated into the peritoneal cavity with a 15 teflon needle-cannula 1 h after administration of the anaesthetic. The amount of fluid injected was determined by weighing the syringe before and after injection of the fluid, using an electronic scale. At the end of each analysis period (2, 4 and 6 h), incisions were made in the rats' abdomens with an acusector and all the fluid present in

the peritoneum was aspirated with a 1 ml syringe. After removing the surface fluid, the intestines were carefully shifted from the abdominal cavity to collect the residual fluid remaining on the dorsal wall. The fluid recovered was placed in a beaker and weighed. The change in weight compared to time 0 represented the amount of fluid recovered from the peritoneal solution injected.

A series of *in-vivo* experiments was conducted according to the experimental model described above in order to evaluate the transport ability of various dialysis fluids containing carnitine.

The data relating to initial and final weight of the fluid recovered from the peritoneum of the animals at the different analysis times were used to calculate the percentage increase in volume in each animal.

Table 2 gives the data for an experiment in which glucose was used at different concentrations (1.5, 2.5 and 4.25%) as the dialysis fluid. These hyperosmolar solutions constitute our control data since they are those commonly used in clinical practice.

TABLE 2.

Percentage changes in fluid volume recovered from rats submitted to
peritoneal dialysis with 15 ml of solution containing different
concentrations of glucose.

Solutions	% increase in volume (2 h)	% increase in volume (4 h)	% increase in volume (6 h)
Glucose 1.5%	21.1±1.5	21.5±1.9	21.7±1.6
Glucose 2.5%	35.2±1.4	35.9±1.3	36.8±1.5
Glucose 4.25%	59.8±1.6	60.5±1.7	59.1±1.6

Results are expressed as mean (n=3) ± S.D.

Glucose at all concentrations used causes an increase in intraperitoneal fluid volume which is completed in the first 2 h. In fact, at 4 and 6 h the fluid volume in the peritoneum remains constant.

The same experiment was conducted using scalar concentrations of carnitine. The results are given in Table 3.

TABLE 3.

Percentage changes in fluid volume recovered from rats submitted to peritoneal dialysis with 15 ml of solution containing different concentrations of carnitine.

5

Solutions	% increase in volume (2 h)	% increase in volume (4 h)	% increase in volume (6 h)
Carnitine 1.5%	25.8±1.3	22.5±1.6	25.3±1.4
Carnitine 2.5%	38.1±1.4	37.3±1.4	38.5±1.6
Carnitine 4.25%	61.5±1.5	64.1±1.8	62.3±1.9

Results are expressed as mean (n=3) ± S.D.

Carnitine, too, proves to be a good osmotic agent, at least as good as glucose. The percentage increase in intraperitoneal fluid volume is slightly greater than that produced by glucose. The fluid recovery by the solution containing carnitine is also rapid in this case and reaches peak activity within 2 h, producing no further increases in volume at the later observation times (4, 6 h).

After confirming the activity of carnitine as an osmotic agent for peritoneal dialysis *in vivo*, we conducted a series of experiments using carnitine in mixtures with glucose or amino acids (aa), maintaining the total percentage of osmolites equal to 1.5%. The results are summarized in Table 4.

./.

Table 5, on the other hand, gives the composition of the amino acids used. The composition of the amino acid solution is the optimal composition to minimize the metabolic acidosis that may occur when such solutions are used.

TABLE 4.

Percentage changes in fluid volume recovered from rats submitted to peritoneal dialysis with 15 ml of solution containing different mixtures of carnitine, glucose and amino acids (aa).

5

Solutions	% increase in volume (2 h)	% increase in volume (4 h)	% increase in volume (6 h)
Carnitine 1.5%	25.8±1.3	22.5±1.6	25.3±1.4
Glucose 1.5%	21.1±1.5	21.5±1.9	21.7±1.6
Car+Glu 1.0+0.5%	25.6±1.6	28.4±1.6	27.6±1.5
Car+aa 0.5+1.0%	24.6±1.2	26.4±1.3	25.3±1.2
Car+aa 0.8+0.7%	23.1±1.5	24.5±1.4	26.0±1.5
Car+aa 1.0+0.5%	25.9±1.5	28.4±1.5	33.8±1.4

Results are expressed as mean (n=3) ± S.D.

The fluid recovery in the first 2 h is comparable for all solutions assayed and ranges from 21.7 to 25.9%. Moreover, this increase remains constant at the later observation times (4 and 6 h) for all solutions, except for a tendency to increase over time in the case of the solution containing Car+aa (1.0+0.5%).

./.

TABLE 5

Composition of solution containing aa

Amino acids	Concentration (mg %)
Leucine	74-112
Valine	100-151
Threonine	47-71
Isoleucine	61-92
Lysine HCl	55-83
Histidine	52-78
Methionine	32-48
Phenylalanine	42-62
Tryptophan	20-30
Alanine	68-103
Proline	43-65
Arginine	60-113
Glycine	36-55
Serine	48-72
Tyrosine	20-35
Aspartate	55-83
Glutamate	55-83
Phenylalanine/Tyrosine	1.3-3.0
Generating/neutralizing acids	1-2.2
Essential/Total	0.4-0.7

5

The following examples further illustrate the invention.

./.

EXAMPLE 1

Solution for peritoneal dialysis

5	Sodium	134.0	mmol/l
	Calcium	1.75	mmol/l
	Magnesium	0.5	mmol/l
	Chloride	103.5	mmol/l
	Bicarbonate	34.0	mmol/l
10	L-carnitine	1.5	%

EXAMPLE 2

Solution for peritoneal dialysis

15	Sodium	134.0	mmol/l
	Calcium	1.75	mmol/l
	Magnesium	0.5	mmol/l
	Chloride	103.5	mmol/l
20	Bicarbonate	34.0	mmol/l
	L-carnitine	2.5	%

EXAMPLE 3

Solution for peritoneal dialysis

25	Sodium	134.0	mmol/l
	Calcium	1.75	mmol/l
	Magnesium	0.5	mmol/l
	Chloride	103.5	mmol/l
30	Bicarbonate	34.0	mmol/l
	L-carnitine	4.25	%

EXAMPLE 4

Solution for peritoneal dialysis

5	Sodium	134.0	mmol/l
	Calcium	1.75	mmol/l
	Magnesium	0.5	mmol/l
	Chloride	103.5	mmol/l
	Lactate	35.0	mmol/l
10	L-carnitine	1.0	%
	Glucose	0.5	%

EXAMPLE 5

Solution for peritoneal dialysis

15	Sodium	134.0	mmol/l
	Calcium	1.75	mmol/l
	Magnesium	0.5	mmol/l
	Chloride	103.5	mmol/l
20	Bicarbonate	34.0	mmol/l
	L-carnitine	1.0	%

Mixture of amino acids as in Table 5 0.5 %

EXAMPLE 6

25 Solution for twin-bag peritoneal analysis

Bag 1

	Sodium	193.0	mmol/l
	Calcium	1.75	mmol/l
	Magnesium	0.5	mmol/l
30	Chloride	103.5	mmol/l
	Lactate	35.0	mmol/l
	Glucose	0.5-4.0	%

Bag 2

	Bicarbonate	34.0	mmol/l
35	L-carnitine	4.0-0.5	%

EXAMPLE 7

Solution for peritoneal dialysis

5	Sodium	134.0	mmol/l
	Potassium	2.0	mmol/l
	Calcium	1.75	mmol/l
	Magnesium	0.5	mmol/l
	Chloride	105.5	mmol/l
10	Bicarbonate	34.0	mmol/l
	L-carnitine	1.5-4.25	%

EXAMPLE 8

Solution for peritoneal dialysis

15	Sodium	134.0	mmol/l
	Potassium	2.0	mmol/l
	Calcium	1.75	mmol/l
	Magnesium	0.5	mmol/l
20	Chloride	105.5	mmol/l
	Lactate	35.0	mmol/l
	L-carnitine	0.5-4.0	%
	Glucose	4.0-0.5	%

CLAIMS

1. Use of L-carnitine and/or its alkanoyl derivatives, in which the alkanoyl is a straight or branched aliphatic group, with 2 to 8 carbon atoms, optionally in the form of a pharmaceutically acceptable salt, as an osmotic agent in solutions for medical use.
2. Use according to claim 1, in which said osmotic agent is L-carnitine.
3. Use according to claim 1, in which said osmotic agent is acetyl L-carnitine.
4. Use according to any of claims 1-3, where said osmotic agent is a combination of L-carnitine and at least one of its alkanoyl derivatives.
5. Use according to claim 4, in which said alkanoyl derivative is acetyl L-carnitine.
6. Use according to any of claims 1-3, or either of claims 4 or 5, in which at least one other osmotic agent is combined in the solution for medical use.
7. Use according to claim 6, in which said osmotic agent is glucose.
8. Use according to any of the foregoing claims, in which said solution for medical use is used for peritoneal dialysis.

9. Solution for medical use characterized in that the osmotic agent is L-carnitine, and/or one or more alkanoyl derivatives of L-carnitine, in which the alkanoyl is a straight or branched aliphatic group with 2 to 8 carbon atoms, optionally in the form of a salt with a pharmaceutically acceptable acid.
10. Solution for peritoneal dialysis characterized in that the osmotic agent is L-carnitine, and/or one or more alkanoyl derivatives of L-carnitine, in which the alkanoyl is a straight or branched aliphatic group with 2 to 8 carbon atoms, optionally in the form of a salt with a pharmaceutically acceptable acid.
11. Solution according to claim 10, in which the osmotic agent is L-carnitine in combination with at least one of its alkanoyl derivatives, in which the alkanoyl is a straight or branched aliphatic group with 2 to 8 carbon atoms, optionally in the form of a salt with a pharmaceutically acceptable acid.
12. Solution according to claim 11, in which said alkanoyl derivative is acetyl L-carnitine, optionally in the form of a salt with a pharmaceutically acceptable acid.
13. Solution according to any of claims 10-12, in which said pharmaceutically acceptable salt is selected from the group consisting of fumarate, aspartate, citrate and maleate,
14. Solution according to any of claims 10-13 in the form of a

concentrate.

15. Solution according to any of claims 10-13, in which said osmotic agent is L-carnitine at 1.5% concentration w/v.
16. Solution according to any of claims 10-14, in which said
5 osmotic agent consists of a combination of L-carnitine and another osmotic agent.
17. Solution for peritoneal dialysis, characterized in that the osmotic agent is L-carnitine at concentrations ranging from approximately 0.5 to approximately 10% w/v.
- 10 18. Solution for peritoneal dialysis, characterized in that the osmotic agent is L-carnitine at concentrations ranging from approximately 0.7 to approximately 7% w/v.
19. Solution for peritoneal dialysis, characterized in that the osmotic agent is L-carnitine at concentrations ranging from
15 approximately 1 to approximately 5% w/v.
20. Solution for peritoneal dialysis, characterized in that the osmotic agent is L-carnitine at a concentration of approximately 1.5% w/v.
21. Solution for peritoneal dialysis, characterized in that the
20 osmotic agent is L-carnitine at a concentration of approximately 2.5% w/v.
22. Solution for peritoneal dialysis, characterized in that the

./.

osmotic agent is L-carnitine at a concentration of approximately 4.25% w/v.

23. Solution for peritoneal dialysis, characterized in that the osmotic agent is a combination of L-carnitine at a concentration of 1.0% w/v and glucose at a concentration of 0.5% w/v.

24. Solution for peritoneal dialysis, characterized in that the osmotic agent is a combination of L-carnitine at a concentration of 0.5% w/v and a mixture of amino acids at a concentration of 1.0% w/v.

25. Solution for peritoneal dialysis, characterized in that the osmotic agent is a combination of L-carnitine at a concentration of 0.8% w/v and a mixture of amino acids at a concentration of 0.7% w/v.

26. Solution for peritoneal dialysis, characterized in that the osmotic agent is a combination of L-carnitine at a concentration of 1.0% w/v and a mixture of amino acids at a concentration of 0.5% w/v.

27. Solution according to any of claims 24-26, in which the mixture of amino acids has the following composition:

Amino acids	Concentration (mg %)
Leucine	74-112
Valine	100-151
Threonine	47-71
Isoleucine	61-92
Lysine HCl	55-83
Histidine	52-78
Methionine	32-48
Phenylalanine	42-62
Tryptophan	20-30
Alanine	68-103
Proline	43-65
Arginine	60-113
Glycine	36-55
Serine	48-72
Tyrosine	20-35
Aspartate	55-83
Glutamate	55-83
Phenylalanine/Tyrosine	1.3-3.0
Generating/neutralizing acid	1-2.2
Essential/Total	0.4-0.7

28. Solution for peritoneal dialysis according to any of the foregoing claims, containing palmitoyl L-carnitine as a surfactant.

29. Container for peritoneal dialysis, containing a solution

5 according to any of the foregoing claims.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IT 99/00317

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/205 A61K31/22 A61P13/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

PAJ, WPI Data, BIOSIS, CHEM ABS Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 059 262 A (SIGMA-TAU) 23 April 1981 (1981-04-23) claims 1-3 page 2, line 89-97 ---	1-12, 14, 17, 18
X	EP 0 627 161 A (SIGMA-TAU) 7 December 1994 (1994-12-07) claims 1-6 page 2, line 58 -page 3, line 11 ---	1-7, 9, 13-23
X	WO 99 07419 A (A. GUPTA) 18 February 1999 (1999-02-18) cited in the application claims 1, 16, 22, 31 page 14, line 27 -page 15, line 12 example 5 --- -/--	1, 2, 8-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 June 2000

Date of mailing of the international search report

30/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Peeters, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IT 99/00317

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 272 549 A (C. CAVAZZA) 9 June 1981 (1981-06-09) cited in the application claims 1,4,6 column 4, line 45-51 ---	1,2,6-10
X	US 4 237 167 A (C. CAVAZZA) 2 December 1980 (1980-12-02) cited in the application claims 1,2 column 4, line 10-18 ---	1,3,6-10
X	B.A. WARADY E.A.: "Carnitine status of pediatric patients on continuous ambulatory peritoneal dialysis" AMERICAN JOURNAL OF NEPHROLOGY, vol. 10, no. 2, 1990, pages 109-114, XP000921067 page 109 page 110, column 1 page 112 -----	1,2, 6-10,16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IT 99/00317

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2059262 A	23-04-1981	IT 1120033 B	19-03-1986
		AT 395111 B	25-09-1992
		AT 495580 A	15-03-1985
		BE 885527 A	02-02-1981
		CA 1154678 A	04-10-1983
		CH 641675 A	15-03-1984
		DE 3037479 A	21-05-1981
		FR 2468371 A	08-05-1981
		IL 61190 A	29-06-1984
		JP 1003847 B	23-01-1989
		JP 1684768 C	31-07-1992
		JP 56061310 A	26-05-1981
		LU 82818 A	02-02-1981
		NL 8005450 A	07-04-1981
		US 4320145 A	16-03-1982
EP 627161 A	07-12-1994	IT 1261695 B	29-05-1996
		AT 170709 T	15-09-1998
		DE 69413120 D	15-10-1998
		DE 69413120 T	11-02-1999
		ES 2122214 T	16-12-1998
		JP 7048329 A	21-02-1995
		US RE36331 E	05-10-1999
		US 5496821 A	05-03-1996
WO 9907419 A	18-02-1999	AU 8898898 A	01-03-1999
		EP 1009452 A	21-06-2000
US 4272549 A	09-06-1981	IT 1156741 B	04-02-1987
		BE 876262 A	03-09-1979
		BE 876263 A	03-09-1979
		CA 1135626 A	16-11-1982
		CA 1134274 A	26-10-1982
		CH 642549 A	30-04-1984
		CH 642550 A	30-04-1984
		DE 2919543 A	22-11-1979
		DE 2919593 A	22-11-1979
		DK 198179 A	16-11-1979
		DK 198279 A	16-11-1979
		FR 2425854 A	14-12-1979
		FR 2425855 A	14-12-1979
		GB 2020552 A,B	21-11-1979
		GB 2020550 A,B	21-11-1979
		IE 48121 B	03-10-1984
		IE 48122 B	03-10-1984
		JP 1834530 C	29-03-1994
		JP 5030809 B	11-05-1993
		JP 54151130 A	28-11-1979
		JP 54151131 A	28-11-1979
		LU 81263 A	10-09-1979
		LU 81264 A	10-09-1979
		NL 7903823 A	19-11-1979
		NL 7903824 A	19-11-1979
		US 4237167 A	02-12-1980
US 4237167 A	02-12-1980	IT 1156741 B	04-02-1987
		BE 876262 A	03-09-1979
		BE 876263 A	03-09-1979

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/IT 99/00317

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4237167 A		CA 1135626 A	16-11-1982
		CA 1134274 A	26-10-1982
		CH 642549 A	30-04-1984
		CH 642550 A	30-04-1984
		DE 2919543 A	22-11-1979
		DE 2919593 A	22-11-1979
		DK 198179 A	16-11-1979
		DK 198279 A	16-11-1979
		FR 2425854 A	14-12-1979
		FR 2425855 A	14-12-1979
		GB 2020552 A,B	21-11-1979
		GB 2020550 A,B	21-11-1979
		IE 48121 B	03-10-1984
		IE 48122 B	03-10-1984
		JP 1834530 C	29-03-1994
		JP 5030809 B	11-05-1993
		JP 54151130 A	28-11-1979
		JP 54151131 A	28-11-1979
		LU 81263 A	10-09-1979
		LU 81264 A	10-09-1979
		NL 7903823 A	19-11-1979
		NL 7903824 A	19-11-1979
		US 4272549 A	09-06-1981